

## References

- [1] A. H. Ali, J. W. P. Hirschfeld and H. Kaneta, The automorphism group of a complete  $(q-1)$ -arc in  $\text{PG}(2, q)$ , *J. Combin. Designs* 2 (1994), 131–145.
- [2] A. H. Ali, J. W. P. Hirschfeld and H. Kaneta, On the size of arcs in projective spaces, *IEEE Trans. Inform. Theory*, 41 (1995), 1649–1656.
- [3] L. R. A. Casse and D. G. Glynn, The solution to Beniamino Segre’s problem  $I_{r,q}$ ,  $r = 3$ ,  $q = 2^h$ , *Geom. Dedicata*, 13 (1982), 157–163.
- [4] J. M. Chao and H. Kaneta, A complete 24-arc in  $\text{PG}(2,29)$  with the automorphism group  $\text{PSL}(2,7)$ , *Rend. Math. Appl.* (7), 16 (1996), 537–544.
- [5] J. M. Chao and H. Kaneta, Classical arcs in  $\text{PG}(r, q)$  for  $11 \leq q \leq 19$ , *Discrete. Math.*, 174 (1997), 87–94.
- [6] J. M. Chao and H. Kaneta, Classical arcs in  $\text{PG}(r, q)$  for  $23 \leq q \leq 29$ , *Discrete. Math.*, 226 (2001), 377–385.
- [7] K. Coolsaet and H. Sticker, A full classification of the complete  $k$ -arcs of  $\text{PG}(2,23)$  and  $\text{PG}(2,25)$ , *Journal of Combinatorial Designs*, 17 (2009), 459–477.
- [8] K. Coolsaet and H. Sticker, A full classification of the complete  $k$ -arcs in  $\text{PG}(2,27)$ , *Designs, Codes and Cryptography*, submitted paper.
- [9] G. Faina, S. Marcugini, A. Milani and F. Pambianco, The spectrum of the values  $k$  for which there exists a complete  $k$ -arc in  $\text{PG}(2, q)$  for  $q \leq 23$ , *Ars Combin.*, 47 (1997) 3–11.
- [10] G. Faina and F. Pambianco, On some 10-arcs for deriving the minimum order for complete arcs in small projective planes, *Discrete Math.*, 208–209 (1999), 261–271.
- [11] J. C. Fisher, J. W. P. Hirschfeld and J. A. Thas, Complete arcs in planes of square order, *Combinatorics '84 Ann. Discrete Math.* 30, North-Holland Mathematics Studies 123, North-Holland, Amsterdam, 1986, 243–250.
- [12] D. G. Glynn, The non-classical 10-arcs of  $\text{PG}(4,9)$ , *Discrete Math.* 59 (1986), 43–51.
- [13] C. E. Gordon, Orbits of arcs in  $\text{PG}(N, K)$  under projectivities, *Geom. Dedicata*, 42 (1992), no. 2, 187–203.

- [14] J. W. P. Hirschfeld, The main conjecture for MDS codes, *Cryptography and Coding*, 5th IMA Conference, Cirencester, UK, 1995, (C. Boyd ed.), Springer, 44–52.
- [15] J. W. P. Hirschfeld, *Projective Geometries over Finite Fields*, 2nd ed., Clarendon, Oxford (1998).
- [16] J. W. P. Hirschfeld and G. Korchmáros, Arcs and curves over a finite field, *Finite Field Appl.*, 5 (1999), 393–408.
- [17] J. W. P. Hirschfeld and L. Storme, The packing problem in statistics, coding theory and finite projective spaces, *Journal of Statistical Planning and Inference*, 72 (1998), 355–380.
- [18] H. Kaneta and T. Maruta, An elementary proof and an extension of Thas’ theorem on  $k$ -arcs, *Math. Proc. Cambridge Philos. Soc.*, 105 (1989), 459–462.
- [19] G. Kéri, Types of superregular matrices and the number of  $n$ -arcs and complete  $n$ -arcs in  $\text{PG}(r, q)$ , *J. Combin. Designs* 14 (2006), 363–390.
- [20] G. Korchmáros, Osservazioni sui risultati di B. Segre relativi a  $k$ -archi contenenti  $k - 1$  punti di un’ovale, *Atti Accad. Naz. Lincei, Rend.* 56 (1974), 541–549.
- [21] S. Marcugini, A. Milani and F. Pambianco, A computer search for complete arcs in  $\text{PG}(2, q)$ ,  $q \leq 128$ , *Rapporto Tecnico n. 18/95*, Università degli Studi di Perugia, 1995.
- [22] S. Marcugini, A. Milani and F. Pambianco, A computer search for small and large complete arcs in  $\text{PG}(2, q)$ , *Rapporto Tecnico n. 5/98*, Università degli Studi di Perugia, 1998.
- [23] S. Marcugini, A. Milani and F. Pambianco, Minimal complete arcs in  $\text{PG}(2, q)$ ,  $q \leq 29$ , *J. Combin. Math. Combin. Comput.*, 47 (2003), 19–29.
- [24] S. Marcugini, A. Milani and F. Pambianco, Complete arcs in  $\text{PG}(2, 25)$ : the spectrum of the sizes and the classification of the smallest complete arcs, *Discrete. Math.*, 307 (2007), 739–747.
- [25] T. Maruta and H. Kaneta, On the uniqueness of  $(q+1)$ -arcs of  $\text{PG}(5, q)$ ,  $q = 2^h$ ,  $h \geq 4$ , *Math. Proc. Cambridge Philos. Soc.* 110 (1991), 91–94.
- [26] F. J. MacWilliams and N. J. A. Sloane, *The Theory of Error-Correcting Codes*, North-Holland, Amsterdam (1977).
- [27] G. Pellegrino, Un’osservazione sul problema dei  $k$ -archi completi in  $S_{2,q}$ , con  $q \equiv 1 \pmod{4}$ , *Atti Accad. Naz. Lincei, Rend.* 63 (1977), 33–44.

- [28] T. Penttila and G. F. Royle, Classification of hyperovals in  $PG(2, 32)$ , *J. Geom.*, 50 (1994), 151–158.
- [29] R.M. Roth and A. Lempel, On MDS codes via Cauchy matrices, *IEEE Trans. Inform. Theory*, 35 (1989), 1314–1319.
- [30] R.M. Roth and G. Seroussi, On generator matrices of MDS codes, *IEEE Trans. Inform. Theory*, 31 (1985), 826–830.
- [31] B. Segre, Curve razionali normali e  $k$ -archi negli spazi finiti, *Ann. Mat. Pura Appl.* 39 (1955), 357–379.
- [32] H. Sticker, Private communication (2007).
- [33] L. Storme and J. A. Thas, Complete  $k$ -arcs in  $PG(n, q)$ ,  $q$  even, *Discrete Math.* 106/107 (1992), 455–469.
- [34] L. Storme and J. A. Thas, MDS codes and arcs in  $PG(n, q)$  with  $q$  even: an improvement of the bounds of Bruen, Thas, and Blokhuis, *J. Combin. Theory Ser. A* 62 (1993), 139–154.
- [35] T. Szőnyi, Arcs, caps, codes and 3-independent subsets, *Giornate di Geometrie Combinatorie (Proc. International Conference Univ. Perugia 1992, G. Faina and G. Tallini, eds.)*, Università degli studi di Perugia, 1993, 57-80.
- [36] J. A. Thas, Complete arcs and algebraic curves in  $PG(2, q)$ , *J. Algebra* 106 (1987), 451–464.
- [37] J. A. Thas, MDS codes and arcs in projective spaces, *Le Matematiche* 47 (1992), 315–328.
- [38] J. A. Thas, Normal rational curves and  $k$ -arcs in Galois spaces, *Rend. Math. Appl.* (6), 1 (1968), 331–334.
- [39] J. F. Voloch, Arcs in projective planes over prime fields, *J. Geom.*, 38 (1990), 198–200.
- [40] J. F. Voloch, Complete arcs in Galois planes of non-square order, *Advances in Finite Geometries and Designs*, Oxford University Press, Oxford, 1991, 401–406.